

09: Quadratic Equations

Key Terms

Equation: a statement that two expressions have the same value.

Quadratic equations: an equation that has the standard form $ax^2 + bx + c = 0$.

Zero product property: if $ab = 0$, the $a = 0$ or $b = 0$.

Factor: to rewrite an expression as a product.

Solutions or roots of the equation: values an equation takes when the values of its domain are substituted for the variable.

Solution set: collection of all solutions to an equation.

Quadratic inequality: a quadratic equation where the equal symbol is replaced by an inequality symbol.

Perfect square trinomial: $a^2 - 2ab + b^2 = (a - b)^2$;
 $a^2 + 2ab + b^2 = (a + b)^2$

Difference of two squares: $a^2 - b^2 = (a - b)(a + b)$

Discriminant: the value under the radical in the quadratic formula, $b^2 - 4ac$.

Quadratic function: function in the form $f(x) = ax^2 + bx + c$, where a, b , and c are real numbers and $a \neq 0$.

Parabola: the graph of a quadratic function.


Axis of symmetry: divides parabola into two equal parts, each part is a mirror image of another.

Vertex: point where the parabola intercepts the axis of symmetry.

x-intercepts: the points where parabolas intercepts x-axis (where $y = 0$).

y-intercepts: point where the parabola intercepts the y-axis (where $x = 0$).

Quadratic Function Graph



A parabola can open downward ($a < 0$) or upward ($a > 0$).

Example: Factoring

Solve. $x^2 - x - 2 = 0$

Solution:

$$\begin{aligned} (x - 2)(x + 1) &= 0 && \text{Factor left side} \\ x - 2 = 0 \text{ or } x + 1 = 0 &&& \text{Apply zero-product} \\ x = 2 \text{ or } x = -1 &&& \text{Solve equations} \end{aligned}$$

The solution set of this equation is $\{-1, 2\}$.

Example: Square Root Method


Solve using the square root method. $x^2 - 9 = 0$

Solution:

$$\begin{aligned} x^2 &= 9 \\ x &= \pm\sqrt{9} \\ x &= 3 \text{ or } x = -3 \end{aligned}$$

The solution set of this equation is $\{-3, 3\}$.

Concept Map



Example: Quadratic Inequality

Find the solution set. $x^2 + 3x < 18$

Solution: Put the equation in standard form.

$$x^2 + 3x - 18 < 0$$

To define boundaries, change the inequality to an equality then find the solution of the equation.


$$x^2 + 3x - 18 = 0$$

$$(x + 6)(x - 3) = 0$$

$$x = -6 \text{ or } 3$$

Denote the test intervals: $(-\infty, -6)$, $(-6, 3)$, $(3, \infty)$.

Find the sign, positive or negative, in each interval using test values.



The solution set of the inequality is the interval $(-6, 3)$.

Example: Quadratic Formula

Solve using the quadratic formula. $2x^2 - 3x + 1 = 0$

Solution: Identify a, b , and c of the quadratic equation, then use the quadratic formula to solve.

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-(-3) \pm \sqrt{1}}{2(2)} \\ &= 1 \text{ and } \frac{1}{2} \end{aligned}$$

The solution set of this equation is $\{\frac{1}{2}, 1\}$.

Example: Complete the Square

Solve by completing the square. $4x^2 - 12x = -5$

Solution:

$$\begin{aligned} 4x^2 - 12x &= -5 \\ 4x^2 - 2(3)(2x) + 9 &= -5 + 9 \\ (2x - 3)^2 &= 4 \end{aligned}$$

Apply the square root method.

$$2x - 3 = \sqrt{4} = 2 \text{ or } 2x - 3 = -\sqrt{4} = -2$$

Solve the equations.

$$\begin{aligned} 2x - 3 &= 2 && 2x - 3 = -2 \\ 2x - 3 + 3 &= 2 + 3 && 2x - 3 + 3 = -2 + 3 \\ 2x &= 5 && \text{or} && 2x = 1 \\ x &= \frac{5}{2} && && x = \frac{1}{2} \end{aligned}$$

The solution set of this equation is $\{\frac{1}{2}, \frac{5}{2}\}$.

How to Use This Cheat Sheet: These are the key concepts related this topic. Try to read through it carefully twice then rewrite it out on a blank sheet of paper. Review it again before the exam.